

Brain tissue segmentation on Diffusion Weighted Magnetic Resonance data

O. Esteban^{1,2}, S. Gorthi¹, G. Wollny², A. Daducci¹, M.J. Ledesma-Carbayo², A. Santos², J.-P. Thiran¹, M. Bach-Cuadra^{3,1}

¹LTS5 (Signal Processing Lab 5), École Polytechnique Fédérale de Lausanne, Lausanne, Switzerland

²BIT (Biomedical Image Technologies), Universidad Politécnica de Madrid - CIBER-BBN, Spain

³Radiology Department, Centre Hospitalier Universitaire Vaudois and University of Lausanne, Lausanne, Switzerland




Motivation


- Application framework: fiber tracking
- DWI clinical practice: cerebral ischemia, epilepsy, MS, etc.
- DWI challenges:
 - ▷ Resolution
 - ▷ Low SNR, eddy currents, motion artifacts, signal loss & warping
- Why DWI segmentation?:
 - ▷ DWI-derived data is widely available
 - ▷ Avoid EPI unwarping
- We propose and developed:
 - ▷ A study of statistical properties of DWI data
 - ▷ One statistical segmentation tool
 - ▷ A validation framework
 - ▷ Comparison to previous works

Methods

- Preliminary study on manually labeled samplings:
 - ▷ PCA: two *families* of maps
 - ▷ LDA: *discriminant* maps
 - ▷ Result: dimensionality & map selection
- Segmentation tool
 - ▷ Multivariate Gaussian Mixture Model (MGMM)
 - ▷ K-Means initialization (ITK based)
 - ▷ E-M optimization (ITK based)
 - ▷ Markov Random Field (MRF) using Graph-Cuts (GC)
 - ▶ α -Expansion or $\alpha\beta$ -Swap available
 - ▶ maxflow-mincut algorithm from [1]
- Validation: fuzzy Jaccard index [2]:



$$TC = \frac{\sum_k \alpha_k \sum_{i \in \mathcal{S}} \min(P(x_i = l_k), P(\hat{x}_i = l_k))}{\sum_k \alpha_k \sum_{i \in \mathcal{S}} \max(P(x_i = l_k), P(\hat{x}_i = l_k))}$$




Conclusions

- Bivariate model with selected maps is reliable and accurate
- Previous works [3, 4] propose the fusion of univariate clusterings
 - ▷ Only for DTI (not DSI, QBI, HARDI, etc.)
 - ▷ No probabilistic results are provided by these methods
- The use of GC speeds up convergence of MRF and increases robustness
- Future lines:
 - ▷ Study the inclusion of PVE and its impact on overall performance and results
 - ▷ Update the framework with more evolved algorithms (variational inference)
 - ▷ Release the code

Dataset

Subjects	Modality	Vol/Subj	Resolution (mm)	Size (voxels)
6	DTI	2 × 63	2.21×2.21×3.00	96×96×34
	T1	1	1.00×1.00×1.20	240×256×260
1	DSI	1 × 515	2.21×2.21×3.00	96×96×44
	T1	1	1.00×1.00×1.20	240×256×260

- DWI-derived scalar maps:
 - ▷ DTI: B0, FA, ADC, AD, TR, PD₁, PD₂, PD₃, RD, λ_1 , λ_2 , λ_3
 - ▷ DSI: B0, gFA, aADC, asADC, afADC, kurtosis, skewness, P0
 - ▷ No fieldmappings were acquired for any case



Results

- PCA & LDA:
 - ▷ Non-hyperspherical distributions
 - ▷ Map selection: FA+ADC (DTI), gFA+aADC (DSI)
 - ▷ LDA functions
- Final benchmarks and comparison
 - ▷ Better than [4] and mFAST
 - ▷ LDA functions useless




Figure : Joint histogram of FA and ADC. In white, iso-contours of the fitted MGMM




Figure : Liu et al.[4]




Figure : mFAST





Figure : Presented tool



Acknowledgements and References

This study is supported by the Center for Biomedical Imaging (CIBM) of the Geneva and Lausanne Universities and the EPFL, as well as the Leenaards and Louis Jeantet foundations. O. Esteban holds a grant from Federal Commission for Scholarships for Foreign Students (ESKAS 2011-12) of the Swiss Government.

References

- [1] Y. Boykov, O. Veksler, and R. Zabih. "Fast approximate energy minimization via graph cuts". In: *Pattern Analysis and Machine Intelligence, IEEE Transactions on* 23.11 (2001), pp. 1222–1239.
- [2] W.R. Crum, O. Camara, and D.L.G. Hill. "Generalized overlap measures for evaluation and validation in medical image analysis". In: *Medical Imaging, IEEE Transactions on* 25.11 (2006), pp. 1451–1461.
- [3] D. Han et al. "An experimental evaluation of diffusion tensor image segmentation using graph-cuts". In: *Engineering in Medicine and Biology Society, 2009. EMBC 2009. Annual International Conference of the IEEE*. 2009, pp. 5653–5656.
- [4] Tianming Liu et al. "Brain tissue segmentation based on DTI data". In: *NeuroImage* 38.1 (Oct. 2007), pp. 114–123. ISSN: 1053-8119. DOI: 10.1016/j.neuroimage.2007.07.002.